

INHIBITORS OF c-JUN N-TERMINAL KINASES (JNK) AND OTHER  
PROTEIN KINASES

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to US  
Provisional Patent Application 60/251,409 filed  
December 5, 2000, the contents of which are  
5 incorporated herein by reference.

TECHNICAL FIELD OF INVENTION

10005133-120501  
The present invention relates to inhibitors  
of protein kinase, especially c-Jun N-terminal kinases  
(JNK), which are members of the mitogen-activated  
protein (MAP) kinase family. There are a number of  
10 different genes and isoforms which encode JNKs.  
Members of the JNK family regulate signal transduction  
in response to environmental stress and proinflammatory  
cytokines and have been implicated to have a role in  
mediating a number of different disorders. The  
15 invention also relates to methods for producing these  
inhibitors. The invention also provides pharmaceutical  
compositions comprising the inhibitors of the invention  
and methods of utilizing those compositions in the  
treatment and prevention of various disorders.

20 BACKGROUND OF THE INVENTION

Mammalian cells respond to extracellular  
stimuli by activating signaling cascades that are  
mediated by members of the mitogen-activated protein  
(MAP) kinase family, which include the extracellular

10005133-120501

signal regulated kinases (ERKs), the p38 MAP kinases and the c-Jun N-terminal kinases (JNKs). MAP kinases (MAPKs) are activated by a variety of signals including growth factors, cytokines, UV radiation, and stress-inducing agents. MAPKs are serine/threonine kinases and their activation occur by dual phosphorylation of threonine and tyrosine at the Thr-X-Tyr segment in the activation loop. MAPKs phosphorylate various substrates including transcription factors, which in turn regulate the expression of specific sets of genes and thus mediate a specific response to the stimulus.

One particularly interesting kinase family are the c-Jun NH<sub>2</sub>-terminal protein kinases, also known as JNKs. Three distinct genes, Jnk1, Jnk2, Jnk3 have been identified and at least ten different splicing isoforms of JNKs exist in mammalian cells [Gupta et al., EMBO J., 15:2760-70 (1996)]. Members of the JNK family are activated by proinflammatory cytokines, such as tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) and interleukin-1  $\beta$  (IL-1 $\beta$ ), as well as by environmental stress, including anisomycin, UV irradiation, hypoxia, and osmotic shock [Minden et al., Biochemica et Biophysica Acta, 1333:F85-F104 (1997)].

The down-stream substrates of JNKs include transcription factors c-Jun, ATF-2, Elk1, p53 and a cell death domain protein (DENN) [Zhang et al. Proc. Natl. Acad. Sci. USA, 95:2586-91 (1998)]. Each JNK isoform binds to these substrates with different affinities, suggesting a regulation of signaling pathways by substrate specificity of different JNKs *in vivo* (Gupta et al., *supra*).

JNKs, along with other MAPKs, have been implicated in having a role in mediating cellular response to cancer, thrombin-induced platelet aggregation, immunodeficiency disorders, autoimmune diseases, cell death, allergies, osteoporosis and heart disease. The therapeutic targets related to activation of the JNK pathway include chronic myelogenous leukemia (CML), rheumatoid arthritis, asthma, osteoarthritis, ischemia, cancer and neurodegenerative diseases.

Several reports have detailed the importance of JNK activation associated with liver disease or episodes of hepatic ischemia [Nat. Genet. 21:326-9 (1999); FEBS Lett. 420:201-4 (1997); J. Clin. Invest. 102:1942-50 (1998); Hepatology 28:1022-30 (1998)]. Therefore, inhibitors of JNK may be useful to treat various hepatic disorders.

A role for JNK in cardiovascular disease such as myocardial infarction or congestive heart failure has also been reported as it has been shown JNK mediates hypertrophic responses to various forms of cardiac stress [Circ. Res. 83:167-78 (1998); Circulation 97:1731-7 (1998); J. Biol. Chem. 272:28050-6 (1997); Circ. Res. 79:162-73 (1996); Circ. Res. 78:947-53 (1996); J. Clin. Invest. 97:508-14 (1996)].

It has been demonstrated that the JNK cascade also plays a role in T-cell activation, including activation of the IL-2 promoter. Thus, inhibitors of JNK may have therapeutic value in altering pathologic immune responses [J. Immunol. 162:3176-87 (1999); Eur. J. Immunol. 28:3867-77 (1998); J. Exp. Med. 186:941-53 (1997); Eur. J. Immunol. 26:989-94 (1996)].

10005133 "120501  
T05021"EE5001

A role for JNK activation in various cancers has also been established, suggesting the potential use of JNK inhibitors in cancer. For example, constitutively activated JNK is associated with HTLV-1 mediated tumorigenesis [Oncogene 13:135-42 (1996)]. JNK may play a role in Kaposi's sarcoma (KS) because it is thought that the proliferative effects of bFGF and OSM on KS cells are mediated by their activation of the JNK signaling pathway [J. Clin. Invest. 99:1798-804 (1997)]. Other proliferative effects of other cytokines implicated in KS proliferation, such as vascular endothelial growth factor (VEGF), IL-6 and TNF $\alpha$ , may also be mediated by JNK. In addition, regulation of the c-jun gene in p210 BCR-ABL transformed cells corresponds with activity of JNK, suggesting a role for JNK inhibitors in the treatment for chronic myelogenous leukemia (CML) [Blood 92:2450-60 (1998)].

JNK1 and JNK2 are widely expressed in a variety of tissues. In contrast, JNK3 is selectively expressed in the brain and to a lesser extent in the heart and testis [Gupta et al., *supra*; Mohit et al., Neuron 14:67-78 (1995); Martin et al., Brain Res. Mol. Brain Res. 35:47-57 (1996)]. JNK3 has been linked to neuronal apoptosis induced by kainic acid, indicating a role of JNK in the pathogenesis of glutamate neurotoxicity. In the adult human brain, JNK3 expression is localized to a subpopulation of pyramidal neurons in the CA1, CA4 and subiculum regions of the hippocampus and layers 3 and 5 of the neocortex [Mohit et al., *supra*]. The CA1 neurons of patients with acute hypoxia showed strong nuclear JNK3-immunoreactivity

10005133.120501  
compared to minimal, diffuse cytoplasmic staining of  
the hippocampal neurons from brain tissues of normal  
patients [Zhang et al., *supra*]. Thus, JNK3 appears to  
be involved involved in hypoxic and ischemic damage of  
5 CA1 neurons in the hippocampus.

In addition, JNK3 co-localizes  
immunochemically with neurons vulnerable in Alzheimer's  
disease [Mohit et al., *supra*]. Disruption of the JNK3  
gene caused resistance of mice to the excitotoxic  
10 glutamate receptor agonist kainic acid, including the  
effects on seizure activity, AP-1 transcriptional  
activity and apoptosis of hippocampal neurons,  
indicating that the JNK3 signaling pathway is a  
critical component in the pathogenesis of glutamate  
15 neurotoxicity [Yang et al., *Nature*, 389:865-870  
(1997)].

Based on these findings, JNK signalling,  
especially that of JNK3, has been implicated in the  
areas of apoptosis-driven neurodegenerative diseases  
20 such as Alzheimer's Disease, Parkinson's Disease, ALS  
(Amyotrophic Lateral Sclerosis), epilepsy and seizures,  
Huntington's Disease, traumatic brain injuries, as well  
as ischemic and hemorrhaging stroke.

Recent evidence now suggests that the closely  
25 related JNK1 and JNK2 isoforms also play a role in  
effecting downstream apoptosis. [Tournier et al.,  
*Science*, 288:870-874 (2000); Mielke et al.,  
*Neuroscience*, 91:471-483 (1999)]. Therefore, these  
isoforms are also potentially important targets for  
30 treating stroke and other apoptosis-related diseases.  
For example, all three JNK isoforms (JNK1, 2 and 3) are  
expressed in the adult brains of humans and rodents.

JNK3 is localized in the hippocampus, as mentioned above, whereas JNK1 and JNK2 are expressed in other compartments of the brain. [Derijard et al., Cell, 76:1025-1037 (1994); Gupta et al., *supra*; Kyriakis et al., Nature, 369:156-160 (1994)] In cells expressing JNK1 and JNK2, disruption of the Jnk1 gene caused a greater decrease in c-Jun kinase activity than disruption of the Jnk2 gene. However, both of these genes had to be knocked out in order to completely protect against stress-induced apoptosis. [Tournier et al., *supra*]. In view of these findings, it would be particularly desirable to find inhibitors that are active against both JNK1 and JNK2 targets.


There remains a high unmet medical need to develop JNK specific inhibitors that are useful in treating the various conditions associated with JNK activation, especially considering the currently available, relatively inadequate treatment options for the majority of these conditions.

Much work has been done to identify and develop drugs that inhibit MAPKs, such as p38 inhibitors. See, e.g., WO 00/31063, WO 98/52937, WO 98/27098 and WO 95/31451. However, to our knowledge, no MAPK inhibitors have been shown to be specifically selective for one or more JNKs versus other related MAPKs. Furthermore, we know of no inhibitors that are active against both JNK1 and JNK2 or against all three of the aforementioned JNK isoforms.

Accordingly, there is still a great need to develop potent inhibitors of JNKs, including JNK1, 2 and 3 inhibitors, that are useful in treating various conditions associated with JNK activation.

## SUMMARY OF THE INVENTION

It has now been found that compounds of this invention and pharmaceutical compositions thereof are effective as inhibitors of c-Jun N-terminal kinases (JNK). These compounds have the general formula I:




I

wherein G is a small group selected from hydrogen or C<sub>1-3</sub> alkyl, Q is pyridine or pyrimidine, and R<sup>1</sup> - R<sup>3</sup> are as defined below. These compounds are selective JNK inhibitors showing good activity against the three isoforms of JNK (JNK1, JNK2 and JNK3) and relatively low activity against p38 kinase. The compounds are therefore useful for treating JNK-mediated diseases, especially neurodegenerative diseases in which all three JNK isoforms are implicated.

## DETAILED DESCRIPTION OF THE INVENTION

The present compounds have the general formula I:




I

wherein:

R<sup>1</sup> is selected from hydrogen, CONH<sub>2</sub>, T<sub>(n)</sub>-R, or T<sub>(n)</sub>-Ar<sup>1</sup>;

R is an aliphatic or substituted aliphatic group;  
 n is zero or one;  
 T is C(=O), CO<sub>2</sub>, CONH, S(O)<sub>2</sub>, S(O)<sub>2</sub>NH, COCH<sub>2</sub> or CH<sub>2</sub>;  
 R<sup>2</sup> is selected from hydrogen, -R, -CH<sub>2</sub>OR, -CH<sub>2</sub>OH, -CH=O,  
 5     -CH<sub>2</sub>SR, -CH<sub>2</sub>S(O)<sub>2</sub>R, -CH<sub>2</sub>(C=O)R, -CH<sub>2</sub>CO<sub>2</sub>R, -CH<sub>2</sub>CO<sub>2</sub>H, -  
       CH<sub>2</sub>CN, -CH<sub>2</sub>NHR, -CH<sub>2</sub>N(R)<sub>2</sub>, -CH=N-OR, -CH=NNHR, -  
       CH=NN(R)<sub>2</sub>, -CH=NNHCOR, -CH=NNHCO<sub>2</sub>R, -CH=NNHSO<sub>2</sub>R, -  
       aryl, -CH<sub>2</sub>(aryl), -CH<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>NHCOR, -CH<sub>2</sub>NHCONHR, -  
       CH<sub>2</sub>NHCON(R)<sub>2</sub>, -CH<sub>2</sub>NRCOR, -CH<sub>2</sub>NHCO<sub>2</sub>R, -CH<sub>2</sub>CONHR, -  
 10    CH<sub>2</sub>CON(R)<sub>2</sub>, -CH<sub>2</sub>SO<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>(heterocyclyl), or -  
       (heterocyclyl);  
 R<sup>3</sup> is selected from hydrogen, -R, hydroxyalkyl,  
       alkoxyalkyl, alkylthioalkyl, aminoalkyl,  
       alkylaminoalkyl, dialkylaminoalkyl, heterocyclyl,  
 15    heterocyclylalkyl, aryl, aralkyl, or aryloxyalkyl;  
 G is hydrogen or C<sub>1-3</sub> alkyl;  
 Q-NH is



wherein the H of Q-NH is optionally replaced by R, COR,  
 20     S(O)<sub>2</sub>R, or CO<sub>2</sub>R;  
 A is N or CH;  
 Ar<sup>1</sup> is aryl, substituted aryl, heterocyclyl or  
       substituted heterocyclyl, wherein Ar<sup>1</sup> is optionally  
       fused to a partially unsaturated or fully unsaturated  
 25    five to seven membered ring containing zero to three  
       heteroatoms;  
 wherein each substitutable carbon atom in Ar<sup>1</sup>, including  
       the fused ring when present, is optionally and  
       independently substituted by halo, R, OR, SR, OH,  
 30    NO<sub>2</sub>, CN, NH<sub>2</sub>, NHR, N(R)<sub>2</sub>, NHCOR, NHCONHR, NHCON(R)<sub>2</sub>,  
       NRCOR, NHCO<sub>2</sub>R, CO<sub>2</sub>R, CO<sub>2</sub>H, COR, CONHR, CON(R)<sub>2</sub>,



S(O)<sub>2</sub>R, SONH<sub>2</sub>, S(O)R, SO<sub>2</sub>NHR, or NHS(O)<sub>2</sub>R, and wherein each saturated carbon in the fused ring is further optionally and independently substituted by =O, =S, =NNHR, =NNR<sub>2</sub>, =N-OR, =NNHCOR, =NNHCO<sub>2</sub>R, =NNHSO<sub>2</sub>R, or

5 =NR; and

wherein each substitutable nitrogen atom in Ar<sup>1</sup> is optionally substituted by R, COR, S(O)<sub>2</sub>R, or CO<sub>2</sub>R.

As used herein, the following definitions shall apply unless otherwise indicated. The term

10 "aliphatic" as used herein means straight chained, branched or cyclic C<sub>1</sub>-C<sub>12</sub> hydrocarbons, preferably one to six carbons, which are completely saturated or which contain one or more units of unsaturation. For example, suitable aliphatic groups include substituted

15 or unsubstituted linear, branched or cyclic alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl. The term "alkyl" and "alkoxy" used alone or as part of a larger moiety refers to both

20 straight and branched chains containing one to twelve carbon atoms. The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety shall include both straight and branched chains containing two to twelve carbon atoms. The terms "haloalkyl", "haloalkenyl" and

25 "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I. The term "heteroatom" means N, O or S and shall include any oxidized form of nitrogen and sulfur, and the

30 quaternized form of any basic nitrogen.

The term "aryl", used alone or as part of a larger moiety as in "aralkyl", refers to aromatic ring

groups having five to fourteen members, such as phenyl, benzyl, 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl, and heterocyclic aromatic groups or heteroaryl groups such as 2-furanyl, 3-furanyl, N-  
5 imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 3-  
10 pyridazinyl, 3-pyridazinyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 5-tetrazolyl, 2-triazolyl, 5-triazolyl, 2-thienyl, or 3-thienyl.

Aryl groups also include fused polycyclic aromatic ring systems in which a carbocyclic aromatic  
15 ring or heteroaryl ring is fused to one or more other rings. Examples include tetrahydronaphthyl, benzimidazolyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, benzodiazepinyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinolinyl,  
20 isoindolyl, acridinyl, benzoisoxazolyl, and the like. Also included within the scope of the term "aryl", as it is used herein, is a group in which one or more carbocyclic aromatic rings and/or heteroaryl rings are fused to a cycloalkyl or non-aromatic heterocyclyl, for  
25 example, indanyl or tetrahydrobenzopyranyl.

The term "heterocyclic ring" or "heterocyclyl" refers to a non-aromatic ring which includes one or more heteroatoms such as nitrogen, oxygen or sulfur in the ring. The ring can be five,  
30 six, seven or eight-membered and/or fused to another ring, such as a cycloalkyl or aromatic ring. Examples include 3-1H-benzimidazol-2-one, 3-1-alkyl-

benzimidazol-2-one, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholino, 3-morpholino, 4-morpholino, 2-thiomorpholino, 3-thiomorpholino, 4-thiomorpholino, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 4-thiazolidinyl, diazolonyl, N-substituted diazolonyl, 1-phthalimidinyl, benzoxane, benzotriazol-1-yl, benzopyrrolidine, benzopiperidine, benzoxolane, benzothiolane, and benzothiane.

A compound of this invention may contain a ring that is fused to a partially saturated or fully unsaturated five to seven membered ring containing zero to three heteroatoms. Such a fused ring may be an aromatic or non-aromatic monocyclic ring, examples of which include the aryl and heterocyclic rings described above.

An aryl group (carbocyclic and heterocyclic) or an aralkyl group, such as benzyl or phenethyl, may contain one or more substituents. Examples of suitable substituents on the unsaturated carbon atom of an aryl group include a halogen, -R, -OR, -OH, -SH, -SR, protected OH (such as acyloxy), phenyl (Ph), substituted Ph, -OPh, substituted -OPh, -NO<sub>2</sub>, -CN, -NH<sub>2</sub>, -NHR, -N(R)<sub>2</sub>, -NHCOR, -NHCONHR, -NHCON(R)<sub>2</sub>, -NRCOR, -NHCO<sub>2</sub>R, -CO<sub>2</sub>R, -CO<sub>2</sub>H, -COR, -CONHR, -CON(R)<sub>2</sub>, -S(O)<sub>2</sub>R, -SONH<sub>2</sub>, -S(O)R, -SO<sub>2</sub>NHR, or -NHS(O)<sub>2</sub>R, where R is an aliphatic group or a substituted aliphatic group.

1005133-120501

An aliphatic group or a non-aromatic heterocyclic ring may contain one or more substituents. Examples of suitable substituents on the saturated carbon of an aliphatic group or of a non-aromatic heterocyclic ring include those listed above for the unsaturated carbon, such as in an aromatic ring, as well as the following: =O, =S, =NNHR, =NNR<sub>2</sub>, =N-OR, =NNHCOR, =NNHCO<sub>2</sub>R, =NNHSO<sub>2</sub>R, or =NR.

A substitutable nitrogen on an aromatic or non-aromatic heterocyclic ring may be optionally substituted. Suitable substituents on the nitrogen include R, COR, S(O)<sub>2</sub>R, and CO<sub>2</sub>R, where R is an aliphatic group or a substituted aliphatic group.

Compounds derived by making isosteric or bioisosteric replacements of carboxylic acid or ester moieties of compounds described herein are within the scope of this invention. Isosteres, which result from the exchange of an atom or group of atoms to create a new compound with similar biological properties to the parent carboxylic acid or ester, are known in the art. The bioisosteric replacement may be physicochemically or topologically based. An example of an isosteric replacement for a carboxylic acid is CONHSO<sub>2</sub>(alkyl) such as CONHSO<sub>2</sub>Me.



It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms or hydrated forms, all such forms of the compounds being within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical

10005133 120501

isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

Preferred compounds of this invention have one or more, and more preferably all, of the following features:

- (a)  $\text{R}^1$  is selected from hydrogen,  $\text{T}_{(n)}\text{-R}$ , or  $\text{T}_{(n)}\text{-Ar}^1$ ;
- (b)  $\text{R}^2$  is selected from hydrogen,  $\text{-R}$ ,  $\text{-CH}_2\text{OR}$ ,  $\text{CH}_2\text{OH}$ ,  $\text{CH}_2(\text{heterocyclyl})$ ,  $\text{-CH}_2(\text{substituted heterocyclyl})$ ,  $\text{-(heterocyclyl)}$ , or  $\text{-(substituted heterocyclyl)}$ ;
- (c)  $\text{R}^3$  is selected from  $\text{-R}$ , heterocyclyl, heterocyclylalkyl, aryl, or aralkyl and/or;
- (d)  $\text{G}$  is hydrogen or methyl.



One embodiment of this invention relates to compounds of formula I wherein Q is a pyrimidine ring, represented by formula II above. Another embodiment relates to compounds of formula I wherein Q is a pyridine ring, represented by formula III above.

Examples of the R<sup>1</sup> group include, but are not limited to, phenyl, cyclohexyl, pyridyl, naphthyl, and quinolinyl, each of which may be optionally substituted. Examples of the R<sup>2</sup> group include, but are not limited to, hydrogen, methyl, alkoxymethyl (such as methoxymethyl, methoxyethoxymethyl and ethoxymethyl), benzyloxymethyl, and heterocyclylmethyl (such as piperidin-1-ylmethyl, morpholin-4-ylmethyl, and tetrahydrofuran-3-ylmethyl), each of which may be optionally substituted. Examples of the R<sup>3</sup> group include, but are not limited to, phenyl and benzyl, each of which may be optionally substituted.

Table 1 below shows examples of compounds of formula II where Q is a pyrimidine.

No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-1	CH <sub>3</sub>	Phenyl	H	Ph
II-2	CH <sub>3</sub>	4-methoxy-phenyl	H	Ph
II-3	CH <sub>3</sub>	3,4-dimethoxy-phenyl	H	Ph
II-4	CH <sub>3</sub>	3,5-dimethoxy-phenyl	H	Ph
II-5	CH <sub>3</sub>	4-cyano-phenyl	H	Ph
II-6	CH <sub>3</sub>	3-fluoro-phenyl	H	Ph
II-7	CH <sub>3</sub>	4-fluoro-phenyl	H	Ph
II-8	CH <sub>3</sub>	4-COCH <sub>3</sub> -phenyl	H	Ph
II-9	CH <sub>3</sub>	4-CONH <sub>2</sub> -phenyl	H	Ph
II-10	CH <sub>3</sub>	4-SCH <sub>3</sub> -phenyl	H	Ph
II-11	CH <sub>3</sub>	3-OCH <sub>3</sub> -phenyl	H	Ph
II-12	CH <sub>3</sub>	3,4,5-trimethoxy-phenyl	H	Ph

No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-13	CH <sub>3</sub>	4-CO <sub>2</sub> CH <sub>3</sub> -phenyl	H	Ph
II-14	CH <sub>3</sub>	4-SO <sub>2</sub> CH <sub>3</sub> -phenyl	H	Ph
II-15	CH <sub>3</sub>	4-CO <sub>2</sub> CH <sub>3</sub> -phenyl	H	Ph
II-16	CH <sub>3</sub>	4-N(CH <sub>3</sub> ) <sub>2</sub> -phenyl	H	Ph
II-17	CH <sub>3</sub>	3-NO <sub>2</sub> -phenyl	H	Ph
II-18	CH <sub>3</sub>	3-NHCOCH <sub>3</sub> -phenyl	H	Ph
II-19	CH <sub>3</sub>	3-NH <sub>2</sub> -phenyl	H	Ph
II-20	CH <sub>3</sub>	4-NO <sub>2</sub> -phenyl	H	Ph
II-21	CH <sub>3</sub>	3-(CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H)-phenyl	H	Ph
II-22	CH <sub>3</sub>	3-(CH <sub>2</sub> CO <sub>2</sub> H)-phenyl	H	Ph
II-23	CH <sub>3</sub>	3-CH <sub>2</sub> OH-phenyl	H	4-CH <sub>3</sub> -Ph
II-24	CH <sub>3</sub>	Phenyl	H	4-OMe-phenyl
II-25	CH <sub>3</sub>	4-methoxy-phenyl	H	4-OMe-phenyl
II-26	CH <sub>3</sub>	3,4-dimethoxy-phenyl	H	4-Cl-Ph
II-27	CH <sub>3</sub>	3,5-dimethoxy-phenyl	H	3,4-Cl <sub>2</sub> -Ph
II-28	CH <sub>3</sub>	4-cyano-phenyl	H	4-F-Ph
II-29	CH <sub>3</sub>	3-fluoro-phenyl	H	4-OMe-phenyl
II-30	CH <sub>3</sub>	4-fluoro-phenyl	H	2,5-Cl <sub>2</sub> -PhPh
II-31	CH <sub>3</sub>	4-COCH <sub>3</sub> -phenyl	H	2,4-F <sub>2</sub> -Ph
II-32	CH <sub>3</sub>	4-CONH <sub>2</sub> -phenyl	H	4-NO <sub>2</sub> -Ph
II-33	CH <sub>3</sub>	4-SCH <sub>3</sub> -phenyl	H	3,5-Cl <sub>2</sub> -Ph
II-34	CH <sub>3</sub>	3-OCH <sub>3</sub> -phenyl	H	3-Cl-Ph
II-35	CH <sub>3</sub>	3,4,5-trimethoxy-phenyl	H	4-OMe-phenyl
II-36	CH <sub>3</sub>	4-CH <sub>3</sub> -phenyl	H	3-OBn-Ph
II-37	CH <sub>3</sub>	cyclohexyl	H	4-OMe-phenyl
II-38	CH <sub>3</sub>	cyclohexyl	H	4-OMe-phenyl
II-39	CH <sub>3</sub>	cyclohexyl	H	4-Cl-Ph
II-40	CH <sub>3</sub>	cyclohexyl	H	3,4-Cl <sub>2</sub> -Ph
II-41	CH <sub>3</sub>	cyclohexyl	H	4-F-Ph
II-42	CH <sub>3</sub>	cyclohexyl	H	4-OMe-phenyl
II-43	CH <sub>3</sub>	cyclohexyl	H	2,5-Cl <sub>2</sub> -Ph
II-44	CH <sub>3</sub>	cyclohexyl	H	2,4-F <sub>2</sub> -Ph
II-45	CH <sub>3</sub>	cyclohexyl	H	4-NO <sub>2</sub> -Ph
II-46	CH <sub>3</sub>	cyclohexyl	H	3,5-Cl <sub>2</sub> -Ph
II-47	CH <sub>3</sub>	cyclohexyl	H	3-Cl-Ph

No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-48	CH <sub>3</sub>	cyclohexyl	H	4-OMe-phenyl
II-49	CH <sub>3</sub>	cyclohexyl	H	3-OBn-Ph
II-50	CH <sub>3</sub>	cyclohexyl	H	-CH <sub>2</sub> Ph
II-51	CH <sub>3</sub>	cyclohexyl	H	Ph
II-52	CH <sub>3</sub>	Phenyl	H	4-OMe-phenyl
II-53	H	4-methoxy-phenyl	H	4-OMe-phenyl
II-54	H	3,4-dimethoxy-phenyl	H	4-Cl-Ph
II-55	H	3,5-dimethoxy-phenyl	H	3,4-Cl <sub>2</sub> -Ph
II-56	H	4-cyano-phenyl	H	4-F-Ph
II-57	H	3-fluoro-phenyl	H	4-OMe-phenyl
II-58	H	4-fluoro-phenyl	H	2,5-Cl <sub>2</sub> -PhPh
II-59	H	4-COCH <sub>3</sub> -phenyl	H	2,4-F <sub>2</sub> -Ph
II-60	H	4-CONH <sub>2</sub> -phenyl	H	4-NO <sub>2</sub> -Ph
II-61	H	4-SCH <sub>3</sub> -phenyl	H	3,5-Cl <sub>2</sub> -Ph
II-62	H	3-OCH <sub>3</sub> -phenyl	H	3-Cl-Ph
II-63	H	3,4,5-trimethoxy-phenyl	H	4-OMe-phenyl
II-64	H	4-CH <sub>3</sub> -phenyl	H	3-OBn-Ph
II-65	H	cyclohexyl	H	benzyl
II-66	H	cyclohexyl	H	4-OMe-phenyl
II-67	H	cyclohexyl	H	phenyl
II-68	H	cyclohexyl	H	3,4-Cl <sub>2</sub> -Ph
II-69	H	cyclohexyl	H	2,4-Cl <sub>2</sub> -Ph
II-70	H	cyclohexyl	H	4-OMe-phenyl
II-71	H	cyclohexyl	H	2,5-Cl <sub>2</sub> -Ph
II-72	H	cyclohexyl	H	2,4-F <sub>2</sub> -Ph
II-73	H	cyclohexyl	H	4-NO <sub>2</sub> -Ph
II-74	H	cyclohexyl	H	3,5-Cl <sub>2</sub> -Ph
II-75	H	cyclohexyl	H	3-Cl-Ph
II-76	H	Phenyl	H	4-OMe-phenyl
II-78	CH <sub>3</sub>	4-methoxy-phenyl	H	-CH <sub>2</sub> Ph
II-79	CH <sub>3</sub>	3,4-dimethoxy-phenyl	H	-CH <sub>2</sub> Ph
II-80	CH <sub>3</sub>	3,5-dimethoxy-phenyl	H	-CH <sub>2</sub> Ph
II-81	CH <sub>3</sub>	4-cyano-phenyl	H	-CH <sub>2</sub> Ph
II-82	CH <sub>3</sub>	3-fluoro-phenyl	H	-CH <sub>2</sub> Ph
II-83	CH <sub>3</sub>	3,4,5-trimethoxy-phenyl	H	-CH <sub>2</sub> Ph




No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-84	CH <sub>3</sub>	3-pyridyl	H	Ph
II-85	CH <sub>3</sub>	4-methoxy-pyrid-3-yl	H	Ph
II-86	CH <sub>3</sub>	2-naphthyl	H	Ph
II-87	CH <sub>3</sub>	Isoquinolin-4-yl	H	Ph
II-88	CH <sub>3</sub>	6-methoxy-naphthalen-2-yl	H	Ph
II-89	CH <sub>3</sub>	Indan-1-on-5-yl	H	Ph
II-90	CH <sub>3</sub>	2-methyl-quinolin-6-yl	H	Ph
II-91	CH <sub>3</sub>	4-methoxy-phenyl	CH <sub>3</sub>	Ph
II-92	CH <sub>3</sub>	3,4-dimethoxy-phenyl	CH <sub>3</sub>	Ph
II-93	CH <sub>3</sub>	3,5-dimethoxy-phenyl	CH <sub>3</sub>	4-OMe-phenyl
II-94	CH <sub>3</sub>	cyclohexyl	CH <sub>3</sub>	4-OMe-phenyl
II-95	CH <sub>3</sub>	cyclohexyl	CH <sub>3</sub>	4-Cl-phenyl
II-96	CH <sub>3</sub>	cyclohexyl	CH <sub>3</sub>	Ph
II-97	CH <sub>3</sub>	4-methoxy-phenyl	CH <sub>3</sub>	-CH <sub>2</sub> Ph
II-98	CH <sub>3</sub>	2-methyl-quinolin-6-yl	CH <sub>3</sub>	-CH <sub>2</sub> Ph
II-99	CH <sub>3</sub>	2-methyl-quinolin-6-yl	CH <sub>3</sub>	-CH <sub>2</sub> Ph
II-100	H	4-F-phenyl	CH <sub>3</sub>	Ph
II-101	H	4-Cl-phenyl	CH <sub>3</sub>	Ph
II-102	H	4-NO <sub>2</sub> -phenyl	CH <sub>3</sub>	Ph
II-103	H	cyclohexyl	CH <sub>3</sub>	2,6-difluoro-phenyl
II-104	H	cyclohexyl	CH <sub>3</sub>	3,5-dichloro-phenyl
II-105	H	cyclohexyl	CH <sub>3</sub>	2,4-dichloro-phenyl
II-106	H	cyclohexyl	CH <sub>3</sub>	Ph
II-107	H	3-Cl-phenyl	CH <sub>3</sub>	Ph
II-108	H	3-(benzyloxy)-phenyl	CH <sub>3</sub>	Ph
II-109	H	phenyl	CH <sub>3</sub>	2,4-difluoro-phenyl
II-110	CH <sub>3</sub>	3-Cl-phenyl	H	phenyl
II-111	H	phenyl	H	2,4-difluoro-phenyl
II-112	H	cyclohexyl	H	phenyl
II-113	H	3-Br-phenyl	CH <sub>3</sub>	phenyl
II-114	H	3-I-phenyl	CH <sub>3</sub>	phenyl
II-115	H	2-chloropyridin-5-yl	CH <sub>3</sub>	phenyl
II-116	H	phenyl	CH <sub>3</sub>	pyridin-2-yl
II-117	H	4-F-phenyl	CH <sub>3</sub>	pyridin-2-yl
II-118	H	4-Cl-phenyl	CH <sub>3</sub>	pyridin-2-yl

No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-119	H	3-Cl-phenyl	CH <sub>3</sub>	pyridin-2-yl
II-120	H	4-NO <sub>2</sub> -phenyl	CH <sub>3</sub>	pyridin-2-yl
II-121	H	3-(benzyloxy)-phenyl	CH <sub>3</sub>	pyridin-2-yl
II-122	H	2,6-difluorophenyl	CH <sub>3</sub>	phenyl
II-123	H	phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-124	H	4-F-phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-125	H	4-Cl-phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-126	H	3-Cl-phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-127	H	4-NO <sub>2</sub> -phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-128	H	3-(benzyloxy)-phenyl	CH <sub>3</sub>	3-Cl-phenyl
II-129	H	naphthalen-2-yl	CH <sub>3</sub>	phenyl
II-130	H	3,4-dimethoxyphenyl	CH <sub>3</sub>	phenyl
II-131	H	phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-132	H	4-F-phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-133	H	4-Cl-phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-134	H	3-Cl-phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-135	H	4-NO <sub>2</sub> -phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-136	H	3-(benzyloxy)-phenyl	CH <sub>3</sub>	6-CH <sub>3</sub> -4-CF <sub>3</sub> -pyridin2-yl
II-137	H	3-F-phenyl	CH <sub>3</sub>	pyridin-2-yl
II-138	H	3-chloro-4-methoxyphenyl	CH <sub>3</sub>	pyridin-2-yl
II-139	H	naphthalen-2-yl	CH <sub>3</sub>	pyridin-2-yl
II-140	H	benzimidazol-2-yl	CH <sub>3</sub>	pyridin-2-yl
II-141	H	3-OH-phenyl	CH <sub>3</sub>	phenyl
II-142	H	3-(benzyloxy)-phenyl	CH <sub>3</sub>	phenyl
II-143	H	4-SO <sub>2</sub> NH <sub>2</sub> -phenyl	CH <sub>3</sub>	phenyl
II-144	H	4-SO <sub>2</sub> NH <sub>2</sub> -phenyl	CH <sub>3</sub>	pyridin-2-yl
II-145	H	3- NH <sub>2</sub> -phenyl	CH <sub>3</sub>	phenyl
II-146	CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy-phenyl	CH <sub>3</sub>	phenyl
II-147	CH <sub>2</sub> CH <sub>3</sub>	3-methoxy-phenyl	CH <sub>3</sub>	4-F-phenyl
II-148	CH <sub>2</sub> CH <sub>3</sub>	3-(benzyloxy)-phenyl	CH <sub>3</sub>	4-F-phenyl
II-149	CH <sub>2</sub> CH <sub>3</sub>	3-Cl-phenyl	CH <sub>3</sub>	4-F-phenyl
II-150	CH <sub>2</sub> CH <sub>3</sub>	3-(benzyloxy)-phenyl	CH <sub>3</sub>	3-NO <sub>2</sub> -phenyl
II-151	CH <sub>2</sub> CH <sub>3</sub>	3-methoxy-phenyl	CH <sub>3</sub>	3-NO <sub>2</sub> -phenyl
II-152	CH <sub>2</sub> CH <sub>3</sub>	3-(benzyloxy)-phenyl	CH <sub>3</sub>	3-CF <sub>3</sub> -phenyl
II-153	CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy-phenyl	CH <sub>3</sub>	3-CF <sub>3</sub> -phenyl

No.	G	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
II-154	CH <sub>2</sub> CH <sub>3</sub>	3-Cl-phenyl	CH <sub>3</sub>	3-CF <sub>3</sub> -phenyl
II-155	CH <sub>2</sub> CH <sub>3</sub>	3-methoxy-phenyl	CH <sub>3</sub>	3-CF <sub>3</sub> -phenyl
II-156	CH <sub>2</sub> CH <sub>3</sub>	3-(benzyloxy)-phenyl	CH <sub>3</sub>	4-NO <sub>2</sub> -phenyl
II-157	CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy-phenyl	CH <sub>3</sub>	4-NO <sub>2</sub> -phenyl
II-158	CH <sub>2</sub> CH <sub>3</sub>	3-Cl-phenyl	CH <sub>3</sub>	4-NO <sub>2</sub> -phenyl
II-159	CH <sub>2</sub> CH <sub>3</sub>	3-methoxy-phenyl	CH <sub>3</sub>	4-NO <sub>2</sub> -phenyl

The compounds of this invention may be prepared in general by methods known to those skilled in the art for analogous compounds, as illustrated by the general schemes below and by the preparative examples that follow.

#### Scheme I




10

Reagents and conditions: a) MeI, NaOH (aqueous); (b) COCl<sub>2</sub>, DMF, CHCl<sub>3</sub>; (c) NH<sub>2</sub>NH<sub>2</sub>, MeOH, 65°C; (d) benzylbromide, CH<sub>3</sub>CN, K<sub>2</sub>CO<sub>3</sub>; (e) Oxone<sup>®</sup>, MeOH/H<sub>2</sub>O (1:1); (f) R<sup>1</sup>NH<sub>2</sub>, DMSO, 85°C or R<sup>1</sup>NH<sub>2</sub>(neat), 100°C; (g) ArNHNH<sub>2</sub>, MeOH, 65°C

Scheme I above shows a synthetic scheme that may be used to prepare compounds of the present invention wherein G and R<sup>2</sup> are each hydrogen. Oxone® is potassium peroxymonosulfate. For illustrative purposes, the steps from compound 3 to prepare compound 5 introduce R<sup>1</sup> as a cyclohexyl group and R<sup>3</sup> as a benzyl group. It will be understood to one skilled in the art that other R<sup>1</sup> and R<sup>3</sup> groups may be introduced in a similar fashion using other R<sup>1</sup> amines and other bromides of R<sup>3</sup> or the like.

#### Scheme II




Reagents and conditions: a) i: LDA, THF, -78 °C, ii: 9, -78°C → 0°C; (b) ArNHNH<sub>2</sub>, MeOH, 100°C; (c) DMF-DMA, 100°C; (d) Oxone®, MeOH/H<sub>2</sub>O(1:1); (e) R<sup>1</sup>NH<sub>2</sub>, DMSO, 80°C or R<sup>1</sup>NH<sub>2</sub>(neat), 100°C

Scheme II above shows a synthetic scheme that may be used to prepare compounds of the present


invention where  $R^2$  is hydrogen and G is other than hydrogen.

Scheme III

5



10




- 15 Reagents and conditions: a) i. LDA, THF, -78°C; ii. BnOCH<sub>2</sub>CON(OCH<sub>3</sub>)CH<sub>3</sub>, -78°C → 0°C; (b) DMF-DMA, 50°C; (c) ArNHNH<sub>2</sub>, EtOH, 80°C; (d) Oxone®, MeOH/H<sub>2</sub>O (1:1); (e)

TMSI,  $\text{CHCl}_3$ ; (f)  $\text{PPh}_3$ ,  $\text{CBr}_4$ ,  $\text{CH}_2\text{Cl}_2$ ; (g)  $\text{RNH}_2$ , DMF; (h)  $\text{R}^1\text{NH}_2$ , DMF,  $40^\circ\text{C}$  or  $\text{R}^1\text{NH}_2(\text{neat})$ ,  $100^\circ\text{C}$ ; (i)  $\text{NaH}$ ,  $\text{R}'\text{OH}$ , THF

Scheme III above shows a synthetic scheme for preparing compounds of this invention where G is hydrogen and  $\text{R}^2$  is other than hydrogen. While the above schemes describe routes to compounds wherein Q is a pyrimidine ring, it will be apparent to one skilled in the art that compounds where Q is a pyridine ring may be prepared in an analogous manner.


Scheme IV



Reagents and conditions: (a) Dimethylformamide dimethyl acetal (DMF-DMA),  $\text{CH}_3\text{CN}$ ,  $80^\circ\text{C}$ ; (b)  $\text{Ph-NHNH}_2$ ,  $\text{MeOH}$ ; (c) Dimethylformamide dimethyl acetal (DMF-DMA),  $\text{CH}_3\text{CN}$ ,  $80^\circ\text{C}$ , 18h; (d)  $\text{RNHC}(=\text{NH})\text{NH}_2$ ,  $\text{CH}_3\text{CN}$ ,  $80^\circ\text{C}$ , 24h.

Scheme IV above shows an alternative route to the preparation of the present compounds wherein the pyrimidine ring is made at the end of the synthesis (step d).

Scheme V



Reagents and conditions: (a) N-chlorosuccinimide,  $(\text{CH}_3)_2\text{S}$   $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$ ; (b)  $(\text{CH}_3\text{CO})_2\text{CH}_2$ , EtOH, NaOH, room temperature; (c) dimethylformamide dimethyl acetal (DMF-DMA), toluene, reflux; (d)  $\text{PhNHC}(=\text{NH})\text{NH}_2$ , NaOMe, MeOH, reflux, 36h (sealed tube).

Scheme V above shows an alternative route for the preparation of the present compounds wherein  $\text{R}^2$  is other than hydrogen. After the reactions and standard work-up conditions, compounds 34 and 36 were used in the next step without purification. Compound 35 was purified by column chromatography and compound 37 was purified by preparative HPLC. While this route is illustrated for compound 37, it will be appreciated by one skilled in the art that this general approach is

applicable to the preparation of compounds having other G and R<sup>1</sup>-R<sup>3</sup> groups.

The compounds of this invention were found to be surprisingly potent inhibitors of all three JNK isoforms (JNK1, JNK2 and JNK3). A significant difference between the present compounds and the most closely related prior art compounds was found to be in the improved inhibition of JNK1 and the selectivity for JNK over p38 activity, as described below.

The activity of the JNK inhibitors of this invention may be assayed *in vitro*, *in vivo* or in a cell line. *In vitro* assays include assays that determine inhibition of either the kinase activity or ATPase activity of activated JNK. For example, see the testing examples described below. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to JNK and may be measured either by radiolabelling the inhibitor prior to binding, isolating the inhibitor/JNK complex and determining the amount of radiolabel bound, or by running a competition experiment where new inhibitors are incubated with JNK bound to known radioligands. One may use any type or isoform of JNK, depending upon which JNK type or isoform is to be inhibited.

The JNK inhibitors or pharmaceutical salts thereof may be formulated into pharmaceutical compositions for administration to animals or humans. These pharmaceutical compositions, which comprise an amount of JNK inhibitor effective to treat or prevent a JNK-mediated condition and a pharmaceutically acceptable carrier, are another embodiment of the present invention.



The term "JNK-mediated condition", as used herein means any disease or other deleterious condition in which JNK is known to play a role. Such conditions include, without limitation, inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, cancer, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin-induced platelet aggregation, and conditions associated with prostaglandin endoperoxidase synthase-2.

Inflammatory diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, acute pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome.

Autoimmune diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, chronic thyroiditis, Graves' disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, or graft vs. host disease.

Destructive bone disorders which may be treated or prevented by the compounds of this invention include, but are not limited to, osteoporosis,

osteoarthritis and multiple myeloma-related bone disorder.

Proliferative diseases which may be treated or prevented by the compounds of this invention  
5 include, but are not limited to, acute myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, multiple myeloma and HTLV-1 mediated tumorigenesis.

Angiogenic disorders which may be treated or  
10 prevented by the compounds of this invention include solid tumors, ocular neovascularization, infantile haemangiomas. Infectious diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, sepsis, septic shock,  
15 and Shigellosis.

Viral diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, acute hepatitis infection (including hepatitis A, hepatitis B and hepatitis C),  
20 HIV infection and CMV retinitis.

Neurodegenerative diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis  
25 (ALS), epilepsy, seizures, Huntington's disease, traumatic brain injury, ischemic and hemorrhaging stroke, cerebral ischemias or neurodegenerative disease, including apoptosis-driven neurodegenerative disease, caused by traumatic injury, acute hypoxia,  
30 ischemia or glutamate neurotoxicity.

"JNK-mediated conditions" also include ischemia/reperfusion in stroke, heart attacks,

myocardial ischemia, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, hepatic ischemia, liver disease, congestive heart failure, pathologic immune responses such as that caused by T cell  
5 activation and thrombin-induced platelet aggregation.

In addition, JNK inhibitors of the instant invention may be capable of inhibiting the expression of inducible pro-inflammatory proteins. Therefore, other "JNK-mediated conditions" which may be treated by  
10 the compounds of this invention include edema, analgesia, fever and pain, such as neuromuscular pain, headache, cancer pain, dental pain and arthritis pain.

The compounds of this invention are also useful as inhibitors of Src-family kinases, especially  
15 Src and Lck. For a general review of these kinases see Thomas and Brugge, *Annu. Rev. Cell Dev. Biol.* (1997) 13, 513; Lawrence and Niu, *Pharmacol. Ther.* (1998) 77, 81; Tatosyan and Mizenina, *Biochemistry (Moscow)* (2000) 65, 49. Accordingly, these compounds are useful for  
20 treating diseases or conditions that are known to be affected by the activity of one or more Src-family kinases. Such diseases or conditions include hypercalcemia, restenosis, hypercalcemia, osteoporosis, osteoarthritis, symptomatic treatment of bone  
25 metastasis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, T-cell mediated hypersensitivity disease, Hashimoto's thyroiditis, Guillain-Barre syndrome, chronic obstructive pulmonary disorder,  
30 contact dermatitis, cancer, Paget's disease, asthma, ischemic or reperfusion injury, allergic disease, atopic dermatitis, and allergic rhinitis. Diseases

that are affected by Src activity, in particular, include hypercalcemia, osteoporosis, osteoarthritis, cancer, symptomatic treatment of bone metastasis, and Paget's disease. Diseases that are affected by Lck activity, in particular, include autoimmune diseases, allergies, rheumatoid arthritis, and leukemia. Compounds of formula II and III wherein R<sup>1</sup> is aryl are especially useful for treating diseases associated with the Src-family kinases, particularly Src or Lck.

In addition to the compounds of this invention, pharmaceutically acceptable derivatives or prodrugs of the compounds of this invention may also be employed in compositions to treat or prevent the above-identified disorders.

A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. Particularly favored derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

Pharmaceutically acceptable prodrugs of the compounds of this invention include, without

limitation, esters, amino acid esters, phosphate esters, metal salts and sulfonate esters.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and  $N^+(C_{1-4} \text{ alkyl})_4$  salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a

sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed

5 are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides.

10 Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions

15 or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other

20 commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of

25 formulation.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions.

30 In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically

added. For oral administration in a capsule form,  
useful diluents include lactose and dried cornstarch.  
When aqueous suspensions are required for oral use, the  
active ingredient is combined with emulsifying and  
5 suspending agents. If desired, certain sweetening,  
flavoring or coloring agents may also be added.

Alternatively, the pharmaceutical  
compositions of this invention may be administered in  
the form of suppositories for rectal administration.

10 These can be prepared by mixing the agent with a  
suitable non-irritating excipient which is solid at  
room temperature but liquid at rectal temperature and  
therefore will melt in the rectum to release the drug.  
Such materials include cocoa butter, beeswax and  
15 polyethylene glycols.

The pharmaceutical compositions of this  
invention may also be administered topically,  
especially when the target of treatment includes areas  
or organs readily accessible by topical application,  
20 including diseases of the eye, the skin, or the lower  
intestinal tract. Suitable topical formulations are  
readily prepared for each of these areas or organs.

Topical application for the lower intestinal  
tract can be effected in a rectal suppository  
25 formulation (see above) or in a suitable enema  
formulation. Topically-transdermal patches may also be  
used.

For topical applications, the pharmaceutical  
compositions may be formulated in a suitable ointment  
30 containing the active component suspended or dissolved  
in one or more carriers. Carriers for topical  
administration of the compounds of this invention



include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

The amount of JNK inhibitor that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the

compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

5                   It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of  
10 administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of inhibitor will also depend upon the particular compound in the composition.

15                   According to another embodiment, the invention provides methods for treating or preventing a JNK-mediated condition comprising the step of administering to a patient one of the above-described pharmaceutical compositions. The term "patient", as  
20 used herein, means an animal, preferably a human.

                  Preferably, that method is used to treat or prevent a condition selected from inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious  
25 diseases, degenerative diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, and thrombin-induced platelet aggregation, or any specific  
30 disease or disorder described above.

                  Depending upon the particular JNK-mediated condition to be treated or prevented, additional drugs,


which are normally administered to treat or prevent that condition, may be administered together with the inhibitors of this invention. For example, chemotherapeutic agents or other anti-proliferative agents may be combined with the JNK inhibitors of this invention to treat proliferative diseases.

Those additional agents may be administered separately, as part of a multiple dosage regimen, from the JNK inhibitor-containing composition.

Alternatively, those agents may be part of a single dosage form, mixed together with the JNK inhibitor in a single composition.

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

Example 1




**4-[1-(2,4-difluoro-phenyl)-1H-pyrazol-4-yl]-2-**

**methanesulfanylmethylpyrimidine.** A 25 mL round-bottom flask equipped with a stirbar was charged with 400 mg (2.04 mmol, 1.00 equivalents) of 2-(2-methanesulfanylmethylpyrimidin-4-yl)-malonaldehyde (obtained according to Brown et al., *Aust. J. Chem.* 33 (10), 2291-2298, 1980), 400 mg of 2,4-difluorophenylhydrazine (2.78 mmol, 1.36


equiv.) and 10 mL of methanol. The resulting solution was stirred at 65 °C for 12 h and then concentrated to a yellow oil which was purified by silica gel chromatography (CH<sub>2</sub>Cl<sub>2</sub>, 15 x 1 cm) to provide 315 mg (1.04 mmol, 50.7% yield) of the product as a yellow foam.

Example 2




4-[1-(2,4-difluoro-phenyl)-1H-pyrazol-4-yl]-2-methanesulfonyl-pyrimidine. To a stirred solution of 315 mg of 4-[1-(2,4-difluoro-phenyl)-1H-pyrazol-4-yl]-2-methanesulfanyl-pyrimidine (1.04 mmol, 1 equiv) in 20 mL of a 1:1 mixture of methanol and water (v:v) was added 500 mg of Oxone<sup>®</sup> as a single portion. The reaction mixture was allowed to stir 12 h at 25 °C and then partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water. The water layer was washed twice with CH<sub>2</sub>Cl<sub>2</sub> and the organic layers were combined and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated to give a yellow solid which was used without further purification. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.805 (1 H, d, J = 5 Hz, Ar), 8.66 (1 H, d, J = 3 Hz, Ar), 8.31 (1 H, s, Ar), 7.90-7.89 (1 H, m, Ar), 7.62 (1 H, d, J = 5 Hz, Ar), 7.07-7.03 (2 H, m, Ar), 3.39 (3 H, s, SCH<sub>3</sub>) ppm.

### Example 3




5 **Cyclohexyl-{4-[1-(2, 4-difluoro-phenyl)-1H-pyrazol-4-yl]-pyrimidin-2-yl}-amine.** A 25 mL round-bottom flask equipped with a stirbar was charged with 265 mg (0.787 mmol, 1 equiv) of 4-[1-(2,4- difluoro-phenyl)-1H-pyrazol-4-yl]-2-methanesulfonyl-pyrimidine dissolved in  
10 5 mL of DMSO. To the stirred solution was added 100  $\mu$ L of cyclohexylamine (86.7 mg, 0.874 mmol, 1.10 equiv). The resulting solution was warmed to 85  $^{\circ}$ C and stirred for 12 h. The orange solution was then partitioned between  $\text{CH}_2\text{Cl}_2$  and water. The aqueous layer was  
15 extracted twice with 20 mL of  $\text{CH}_2\text{Cl}_2$  and the organic layers were combined, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to yellow oil, which was purified by silica gel chromatography (1:1  $\text{CH}_2\text{Cl}_2$ :EtOAc,  $R_f$  = 0.2) to yield 35 mg (0.098 mmol, 12.5% yield) of product as a foam.  $^1\text{H}$   
20 NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.37 (1 H, d,  $J$  = 3 Hz, Ar), 8.17 (1 H, d,  $J$  = 5 Hz, Ar), 8.11 (1 H, s, Ar), 7.83-7.79 (1 H, m, Ar), 6.97-6.94 (2 H, m, Ar), 6.64 (1 H,  $J$  = 5 Hz, Ar), 5.04 (1 H, br s, NH), 3.82-3.80 (1 H, m, cy-Hx), 2.01-1.97 (3 H, m, cy-Hx), 1.70-1.67 (2H, m, cy-Hx), 1.59-1.57 (1 H, m, cy-Hx), 1.37-1.35 (2 H, m, cy-Hx), 1.20-1.16 (2 H, m, cy-Hx) ppm. MS (M+H): 356.15.

Example 4




5 1-(2-Methylsulfonyl-pyrimidin-4-yl)-propan-2-one. To a  
stirred solution of (5.23 g, 38.0 mmol) and  
phenanthroline (10 mg) in THF (100 mL) at -78 °C was  
added dropwise a solution of lithium diisopropylamine  
10 (1.5 M in cyclohexane) until all water was destroyed  
and the solution turned dark (~1 mL). Then, additional  
lithium diisopropylamine (30.4 mL) was added slowly  
over 5 min. After 15 min. at -78 °C, the solution was  
allowed to warm to 0 °C and after 15 min recooled to -78  
°C. Then, a solution of N-methoxy-N-methyl-acetamide  
15 (4.72 mL, 45.6 mmol) in THF (15 mL), which was dried  
over 4 molecular sieves for 1 h, was added dropwise  
via dropfunnel over 20 min. After 25 min. at -78 °C,  
the solution was warmed to 0 °C and stirred for 30 min.  
The solution was poured into dilute HCl (1 M, 100 mL),  
20 extracted with Et<sub>2</sub>O (3 × 200 mL), dried (MgSO<sub>4</sub>),  
filtered and concentrated. Flash chromatography (SiO<sub>2</sub>,  
20% EtOAc-hexanes) provided the product (4.75 g, 26.1  
mmol, 69% yield).

Example 5




N-[1-Methyl-2-(2-methylsulfanyl-pyrimidin-4-yl)-  
 ethylidene]-N'-phenyl-hydrazine A stirred solution of  
 1-(2-methylsulfanyl-pyrimidin-4-yl)-propan-2-one (182  
 5 mg, 1.00 mmol) and phenylhydrazine (99  $\mu$ L, 1.00 mmol)  
 in dry EtOH (2 mL) in a sealed tube was heated to 100  $^{\circ}$ C  
 for 15 min. The reaction was concentrated *in vacuo* to  
 yield the crude product as a yellow paste.



4-(3-Methyl-1-phenyl-1H-pyrazol-4-yl)-2-methylsulfanyl-  
 pyrimidine. To the crude paste prepared above was  
 added dimethylformammdide-dimethoxyacetal (DMF-DMA, 1.33  
 mL, 10 mmol). The resulting solution was heated to 100  
 15  $^{\circ}$ C in a sealed tube for three days. The solution was  
 cooled and poured into H<sub>2</sub>O (15 mL) and extracted with  
 EtOAc (3  $\times$  15 mL), The combined organic extracts was  
 washed with saturated NH<sub>4</sub>Cl (15 mL), brine (15 mL),  
 dried (MgSO<sub>4</sub>), filtered and concentrated. Flash  
 20 chromatography (SiO<sub>2</sub>, 30% EtOAc-hexanes) provided the  
 product (225 mg, 0.800 mmol, 80%).


#### Example 6



2-Methanesulfonyl-4-(3-methyl-1-phenyl-1H-pyrazol-4-yl)-pyrimidine To a stirred solution of 4-(3-Methyl-1-phenyl-1H-pyrazol-4-yl)-2-methylsulfany-pyrimidine (225 mg, 0.800 mmol) in MeOH (8 mL) at ambient


5 temperature was added dropwise a solution Oxone® (1.48 g, 2.40 mmol) in H<sub>2</sub>O (8 mL) over 20 min. After 16 h., the solution was poured into H<sub>2</sub>O (50 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 × 25 mL). The combined extracts were dried (MgSO<sub>4</sub>), filtered and concentrated to provide the  
10 product (239 mg, 0.760 mmol, 95%).

Example 7



15 [4-(3-methyl-1-phenyl-1H-pyrazol-4-yl)-pyrimidin-2-yl]-phenylamine A mixture of 2-methanesulfonyl-4-(3-methyl-1-phenyl-1H-pyrazol-4-yl)-pyrimidine (38 mg, 0.12 mmol) and aniline (neat, 100 µL) was heated to 100 °C for 15h. The volatile components were evaporated  
20 under a stream of nitrogen upon warming. Recrystallization from methanol provided the product (23 mg, 0.070 mmol) in 58% yield.

Example 8






10005133-120501  
105021-EE50001

4-guanidino-benzenesulfonamide 4-Amino-benzenesulfonamide (30mmol, 1 equiv.), cyanamide (1.3g, 31mmol, 1.03 equiv.), and 4N hydrogen chloride in dioxane (8mL, 32mmol) was stirred for 10 minutes at room temperature and heated to 80 C for 18 hours. The mixture was diluted with water (30mL) and diethyl ether (50mL). The aqueous layer was washed with ether (30mL) and the organic layers were discarded. The aqueous layer was neutralized with 6N aqueous HCl (6mL) and diluted with ethyl acetate (50mL). The aqueous layer was extracted with ethyl acetate (50mL) four times. The combined organic layers were concentrated under reduced pressure to afford a solid compound that was washed with diethyl ether (30mL) to afford pale yellow title compound. The compound was characterized by LC/MS and HPLC.


Example 9



20

3-Dimethylamino-1-(5-methyl-1-phenyl-1H-pyrazol-4-yl)-propenone 1-(5-Methyl-1-phenyl-1H-pyrazol-4-yl)-ethanone (10mmol) was dissolved in acetonitrile (5mL) and DMF-DMA (2mL) and refluxed for 18 hours. Solvent was removed and triturated with ether (20mL) and the yellow solid was filtered and washed with ether (10mL) to provide the title compound.

### Example 10



5 4-[4-(5-Methyl-1-phenyl-1H-pyrazol-4-yl)-pyrimidin-2-ylamino]-benzenesulfonamide 3-Dimethylamino-1-(5-methyl-1-phenyl-1H-pyrazol-4-yl)-propenone (0.2mmol) and 4-guanidino-benzenesulfonamide (0.2mmol) in acetonitrile (0.25mL) was heated to reflux for 24 hours  
10 and the resulting mixture was diluted with methanol (3mL). The slurry was filtered and the solid was washed with methanol (2mL) and dried under nitrogen to provide the title compound. M+H (407.0); HPLC Rt = 3.168. HPLC analysis was performed with an HP1100 HPLC  
15 and the eight minute analysis method was performed as follows: (0 min, 10%B; 1 min, 30%B; 5 min, 90%B; 6.5min, 90%B; 6.7min 10%B), where A is 0.01% TFA in water and B is 0.01% TFA in acetonitrile. The analytical column was YMC ODS-AQ, 30x100mm with 1mL/min  
20 flow rate using a diode ray detector (5 wavelenths monitored at 300, 280, 254, 220, and 210nm) at 25°C.

### Cloning, Expression and Purification of JNK3 Protein

A BLAST search of the EST database using the  
25 published JNK3α1 cDNA as a query identified an EST clone (#632588) that contained the entire coding sequence for human JNK3α1. Polymerase chain reactions

(PCR) using *pfu* polymerase (Stratagene) were used to introduce restriction sites into the cDNA for cloning into the pET-15B expression vector at the NcoI and BamHI sites. The protein was expressed in *E. coli*.

5 Due to the poor solubility of the expressed full-length protein (Met 1-Gln 422), an N-terminally truncated protein starting at Ser residue at position 40 (Ser 40) was produced. This truncation corresponds to Ser 2 of JNK1 and JNK2 proteins, and is preceded by a methionine  
10 (initiation) and a glycine residue. The glycine residue was added in order to introduce an NcoI site for cloning into the expression vector. In addition, systematic C-terminal truncations were performed by PCR to identify a construct that give rise to diffraction-  
15 quality crystals. One such construct encodes amino acid residues Ser40-Glu402 of JNK3 $\alpha$ 1 and is preceded by Met and Gly residues.

The construct was prepared by PCR using deoxyoligonucleotides:

20 5' GCTCTAGAGCTCCATGGGCAGCAAAAGCAAAGTTGACAA 3' (forward primer with initiation codon underlined) (SEQ ID NO:1) and

5' TAGCGGATCCTCATTCTGAATTCATTACTTCCTTGTA 3' (reverse primer with stop codon underlined) (SEQ ID NO:2) as  
25 primers and was confirmed by DNA sequencing. Control experiments indicated that the truncated JNK3 protein had an equivalent kinase activity towards myelin basic protein when activated with an upstream kinase MKK7 *in vitro*.

30 *E. coli* strain BL21 (DE3) (Novagen) was transformed with the JNK3 expression construct and grown at 30°C in LB supplemented with 100  $\mu$ g/ml

carbenicillin in shaker flasks until the cells were in log phase ( $OD_{600} \sim 0.8$ ). Isopropylthio- $\beta$ -D-galactosidase (IPTG) was added to a final concentration of 0.8 mM and the cells were harvested 2 hours later by

5 centrifugation.

*E. coli* cell paste containing JNK3 was resuspended in 10 volumes/g lysis buffer (50 mM HEPES, pH 7.2, containing 10% glycerol (v/v), 100 mM NaCl, 2 mM DTT, 0.1 mM PMSF, 2  $\mu$ g/ml Pepstatin, 1 $\mu$ g/ml each of

10 E-64 and Leupeptin). Cells were lysed on ice using a microfluidizer and centrifuged at 100,000 x *g* for 30 min at 4 °C. The 100,000 x *g* supernatant was diluted 1:5 with Buffer A (20 mM HEPES, pH 7.0, 10% glycerol (v/v), 2 mM DTT) and purified by SP-Sepharose

15 (Pharmacia) cation-exchange chromatography (column dimensions: 2.6 x 20 cm) at 4 °C. The resin was washed with 5 column volumes of Buffer A, followed by 5 column volumes of Buffer A containing 50 mM NaCl. Bound JNK3 was eluted with a 7.5 column volume linear gradient of

20 50-300 mM NaCl. JNK3 eluted between 150-200 mM NaCl.

#### Activation of JNK3

Five mg of JNK3 was diluted to 0.5 mg/ml in 50 mM HEPES buffer, pH 7.5, containing 100 mM NaCl, 5 mM DTT, 20 mM  $MgCl_2$  and 1 mM ATP. GST-MKK7(DD) was

25 added at a molar ratio of 1:2.5 GST-MKK7:JNK3. After incubation for 30 minutes at 25°C, the reaction mixture was concentrated 5-fold by ultrafiltration in a Centriprep-30 (Amicon, Beverly, MA), diluted to 10 ml and an additional 1 mM ATP added. This procedure was

30 repeated three times to remove ADP and replenish ATP. The final addition of ATP was 5 mM and the mixture incubated overnight at 4°C.

10005133 120501  
10502T EST5000T

The activated JNK3/GST-MKK7(DD) reaction mixture was exchanged into 50 mM HEPES buffer, pH 7.5, containing 5 mM DTT and 5% glycerol (w/v) by dialysis or ultrafiltration. The reaction mixture was adjusted to 1.1 M potassium phosphate, pH 7.5, and purified by hydrophobic interaction chromatography (at 25 °C) using a Rainin Hydropore column. GST-MKK7 and unactivated JNK3 do not bind under these conditions such that when a 1.1 to 0.05 M potassium phosphate gradient is developed over 60 minutes at a flow rate of 1 ml/minute, doubly phosphorylated JNK3 is separated from singly phosphorylated JNK. Activated JNK3 (i.e. doubly phosphorylated JNK3) was stored at -70°C at 0.25-1 mg/ml.

#### 15                    JNK Inhibition Assays

Compounds were assayed for the inhibition of JNK3 by a spectrophotometric coupled-enzyme assay. In this assay, a fixed concentration of activated JNK3 (10 nM) was incubated with various concentrations of a potential inhibitor dissolved in DMSO for 10 minutes at 30°C in a buffer containing 0.1 M HEPES buffer, pH 7.5, containing 10 mM MgCl<sub>2</sub>, 2.5 mM phosphoenolpyruvate, 200 µM NADH, 30 µg/mL pyruvate kinase, 10 µg/mL lactate dehydrogenase, and 200 µM EGF receptor peptide. The EGF receptor peptide has the sequence KRELVEPLTPSGEAPNQALLR, and is a phosphoryl acceptor in the JNK3-catalyzed kinase reaction. The reaction was initiated by the addition of 10 µM ATP and the assay plate is inserted into the spectrophotometer's assay plate compartment that was maintained at 30°C. The decrease of absorbance at 340 nm was monitored as a function of time. The rate data as a function of

inhibitor concentration was fitted to competitive inhibition kinetic model to determine the  $K_i$ .

The JNK1 and JNK2 assays were performed in an analogous manner.

5

#### Cloning of p38 Kinase in Insect Cells

Two splice variants of human p38 kinase, CSBP1 and CSBP2, have been identified. Specific oligonucleotide primers were used to amplify the coding  
10 region of CSBP2 cDNA using a HeLa cell library (Stratagene) as a template. The polymerase chain reaction product was cloned into the pET-15b vector (Novagen). The baculovirus transfer vector, pVL-(His)6-p38 was constructed by subcloning a XbaI-BamHI  
15 fragment of pET15b-(His)6-p38 into the complementary sites in plasmid pVL1392 (Pharmingen).

The plasmid pVL-(His)6-p38 directed the synthesis of a recombinant protein consisting of a 23-residue peptide (MGSSHHHHHHSSGLVPRGSHMLE, where LVPRGS  
20 represents a thrombin cleavage site) fused in frame to the N-terminus of p38, as confirmed by DNA sequencing and by N-terminal sequencing of the expressed protein. Monolayer culture of Spodoptera frugiperda (Sf9) insect cells (ATCC) was maintained in TNM-FH medium (Gibco  
25 BRL) supplemented with 10% fetal bovine serum in a T-flask at 27°C. Sf9 cells in log phase were co-transfected with linear viral DNA of Autographa californica nuclear polyhedrosis virus (Pharmingen) and transfer vector pVL-(His)6-p38 using Lipofectin  
30 (Invitrogen). The individual recombinant baculovirus clones were purified by plaque assay using 1% low melting agarose.

Expression and Purification of Recombinant p38 Kinase

Trichoplusia ni (Tn-368) High-Five™ cells (Invitrogen) were grown in suspension in Excel-405 protein free medium (JRH Bioscience) in a shaker flask at 27°C. Cells at a density of  $1.5 \times 10^6$  cells/ml were infected with the recombinant baculovirus described above at a multiplicity of infection of 5. The expression level of recombinant p38 was monitored by immunoblotting using a rabbit anti-p38 antibody (Santa Cruz Biotechnology). The cell mass was harvested 72 hours after infection when the expression level of p38 reached its maximum.

Frozen cell paste from cells expressing the (His)<sub>6</sub>-tagged p38 was thawed in 5 volumes of Buffer A (50 mM NaH<sub>2</sub>PO<sub>4</sub> pH 8.0, 200 mM NaCl, 2mM β-Mercaptoethanol, 10% Glycerol and 0.2 mM PMSF). After mechanical disruption of the cells in a microfluidizer, the lysate was centrifuged at 30,000 x g for 30 minutes. The supernatant was incubated batchwise for 3-5 hours at 4°C with Talon™ (Clontech) metal affinity resin at a ratio of 1 ml of resin per 2-4 mgs of expected p38. The resin was settled by centrifugation at 500 x g for 5 minutes and gently washed batchwise with Buffer A. The resin was slurried and poured into a column (approx. 2.6 x 5.0 cm) and washed with Buffer A + 5 mM imidazole.

The (His)<sub>6</sub>-p38 was eluted with Buffer A + 100 mM imidazole and subsequently dialyzed overnight at 4°C against 2 liters of Buffer B, (50 mM HEPES, pH 7.5, 25 mM β-glycerophosphate, 5% glycerol, 2mM DTT). The His<sub>6</sub> tag was removed by addition of at 1.5 units thrombin

10005133-120501  
105021-ET50001  
(Calbiochem) per mg of p38 and incubation at 20°C for 2-3 hours. The thrombin was quenched by addition of 0.2 mM PMSF and then the entire sample was loaded onto a 2 ml benzamidine agarose (American International

5 Chemical) column.

The flow through fraction was directly loaded onto a 2.6 x 5.0 cm Q-Sepharose (Pharmacia) column previously equilibrated in Buffer B + 0.2 mM PMSF. The p38 was eluted with a 20 column volume linear gradient  
10 to 0.6M NaCl in Buffer B. The eluted protein peak was pooled and dialyzed overnight at 4°C vs. Buffer C (50 mM HEPES pH 7.5, 5% glycerol, 50 mM NaCl, 2 mM DTT, 0.2 mM PMSF).

The dialyzed protein was concentrated in a  
15 Centriprep (Amicon) to 3-4 ml and applied to a 2.6 x 100 cm Sephacryl S-100HR (Pharmacia) column. The protein was eluted at a flow rate of 35 ml/hr. The main peak was pooled, adjusted to 20 mM DTT, concentrated to 10-80 mgs/ml and frozen in aliquots at  
20 -70°C or used immediately.

#### Activation of p38

p38 was activated by combining 0.5 mg/ml p38 with 0.005 mg/ml DD-double mutant MKK6 in Buffer B +  
25 10mM MgCl<sub>2</sub>, 2mM ATP, 0.2mM Na<sub>2</sub>VO<sub>4</sub> for 30 minutes at 20°C. The activation mixture was then loaded onto a 1.0 x 10 cm MonoQ column (Pharmacia) and eluted with a linear 20 column volume gradient to 1.0 M NaCl in Buffer B. The activated p38 eluted after the ADP and  
30 ATP. The activated p38 peak was pooled and dialyzed against buffer B + 0.2mM Na<sub>2</sub>VO<sub>4</sub> to remove the NaCl. The dialyzed protein was adjusted to 1.1M potassium



phosphate by addition of a 4.0M stock solution and loaded onto a 1.0 x 10 cm HIC (Rainin Hydropore) column previously equilibrated in Buffer D (10% glycerol, 20mM  $\beta$ -glycerophosphate, 2.0mM DTT) + 1.1MK<sub>2</sub>HPO<sub>4</sub>. The protein was eluted with a 20 column volume linear gradient to Buffer D + 50mM K<sub>2</sub>HPO<sub>4</sub>. The double phosphorylated p38 eluted as the main peak and was pooled for dialysis against Buffer B + 0.2mM Na<sub>2</sub>VO<sub>4</sub>. The activated p38 was stored at -70°C.

10

#### p38 Inhibition Assays

##### A. Inhibition of Phosphorylation of EGF Receptor Peptide

This assay is carried out in the presence of 10 mM MgCl<sub>2</sub> and 100 mM HEPES buffer at pH 7.6. For a typical IC<sub>50</sub> determination, a stock solution is prepared containing all of the above components and activated p38 (5 nM). The stock solution is aliquotted into vials. A fixed volume of DMSO or inhibitor in DMSO (final concentration of DMSO in reaction is 5%) is introduced to each vial, mixed and incubated for 15 minutes at room temperature. EGF receptor peptide, KRELVEPLTPSGEAPNQALLR, a phosphoryl acceptor in p38-catalyzed kinase reaction, is added to each vial to a final concentration of 200  $\mu$ M. The kinase reaction is initiated with ATP (100  $\mu$ M) and the vials are incubated at 30°C. After 30 minutes, the reactions are quenched with equal volume of 10% trifluoroacetic acid (TFA).

The phosphorylated peptide is quantified by HPLC analysis. Separation of phosphorylated peptide from the unphosphorylated peptide is achieved on a reverse phase column (Deltapak, 5  $\mu$ m, C18 100D, part

no. 011795) with a binary gradient of water and  
acetonitrile, each containing 0.1% TFA. IC<sub>50</sub>  
(concentration of inhibitor yielding 50% inhibition) is  
determined by plotting the % activity remaining against  
5 inhibitor concentration.

#### B. Inhibition of ATPase Activity

This assay is carried out in the presence of  
10 mM MgCl<sub>2</sub>, 25 mM β-glycerophosphate, 10% glycerol and  
100 mM HEPES buffer at pH 7.6. For a typical Ki  
10 determination, the Km for ATP in the ATPase activity of  
activated p38 reaction is determined in the absence of  
inhibitor and in the presence of two concentrations of  
inhibitor. Ki is determined from the rate data as a  
function of inhibitor and ATP concentrations. A stock  
15 solution is prepared containing all of the above  
components and activated p38 (60 nM). The stock  
solution is aliquotted into vials. A fixed volume of  
DMSO or inhibitor in DMSO (final concentration of DMSO  
in reaction is 2.5%) is introduced to each vial, mixed  
20 and incubated for 15 minutes at room temperature. The  
reaction is initiated by adding various concentrations  
of ATP and then incubated at 30°C. After 30 minutes,  
the reactions are quenched with 50 μl of EDTA (0.1 M,  
final concentration), pH 8.0. The product of p38  
25 ATPase activity, ADP, is quantified by HPLC analysis.

Separation of ADP from ATP is achieved on a  
reversed phase column (Supelcosil, LC-18, 3 μm, part  
no. 5-8985) using a binary solvent gradient of  
following composition: Solvent A - 0.1 M phosphate  
30 buffer containing 8 mM tetrabutylammonium hydrogen

sulfate (Sigma Chemical Co., catalogue no. T-7158),  
Solvent B - Solvent A with 30% methanol.

Table 2 shows the activities of selected  
compounds in the JNK and p38 inhibition assays.

- 5 Compounds having a  $K_i$  less than 2.0 micromolar are rated  
"A", compounds having a  $K_i$  between 2.0 and 10 micromolar  
are rated "B" and compounds having a  $K_i$  greater than 10  
micromolar are rated "C". The last column shows a  
ratio of p38 to JNK3 activity as a measure of the  
10 selectivity of the present compounds.

Table 2. Comparative Activities in the Inhibition  
Assays.

No.	JNK1	JNK2	JNK3	p38	p38/JNK3
II-65	A	A	A	C	>10
II-67	A	A	A	C	>10
II-70	B	B	A	C	≥10
II-71	A	A	A	C	>10
II-72	A	A	A	C	>10
II-100	A	A	A	C	>10
II-101	A	A	A	C	>10
II-102	A	A	A	C	>10

- 15 As can be seen from Table 2, the compounds of  
this invention have good activity against the three  
isoforms of JNK (JNK1, JNK2 and JNK3). The present  
compounds are much less active against p38 and provide  
good selectivity for JNK, as can be seen from the  
20 p38/JNK3 ratio. For typical compounds the inhibitory  
activity against JNK is estimated to be up to two or

three orders of magnitude greater than the inhibitory activity against p38. By contrast, compounds that have larger G substituents (for example, where G is phenyl or 4-fluoro-phenyl) will be much more active against  
5 p38 and have poorer selectivity for the JNK enzyme.

While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments which utilize the compounds and methods of  
10 this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments which have been represented by way of example.

10005133-120501